

Country Situation

Background Statistics

HIV prevalence - adults (ages 15-49) ^[1] *	3.4% [3.1-3.9%]	2009
HIV prevalence - pregnant women (all ages) ^[2] *	3.0%	2009
Number of women living with HIV delivering ^[3]	16,100	2011
Est. # children (ages 0-14) living with HIV ^[1] *	63,000 [32,000-91,000]	2009
Maternal mortality ratio ^[4]	400/100,000	2010
Est. annual births ^[5]	673,000	2010
Infant mortality rate ^[6]	86/1,000	2010
Under-5 mortality rate ^[7]	120/1,000	2010

HIV prevalence among adults (ages 15-49) in Côte d'Ivoire has declined slightly between 2001 and 2009, from 6.5% to 3.4%^[1]. HIV prevalence among pregnant women (all ages) is slightly lower (3.0%) than the general adult population. In 2011, an estimated 16,100 pregnant women living with HIV delivered children and there were an estimated 4,400 new paediatric HIV infections^[3].

Overall, the uptake of antenatal care and availability of PMTCT services in Côte d'Ivoire is moderate. In 2010, approximately 60% of pregnant women attended at least one ANC visit according to facility data^[8] and PMTCT services were available in 52% of ANC facilities^[9]. HIV testing among pregnant women increased from 6% in 2005 to 59% in 2010^[10]. Although 68% of pregnant women living with HIV received more efficacious ARV regimens for PMTCT in 2011^[3], only 44% of HIV-exposed infants received ARVs for PMTCT in 2010^[10].

Côte d'Ivoire has adopted WHO PMTCT Guidelines Option B and a costed national PMTCT scale-up plan (2012-2016) is in place^[11].

Reaching High Level Targets

Global 2015 Targets


- Reduce the number of new HIV infections among children by 90%
- Reduce the number of HIV-associated deaths to women during pregnancy, delivery or puerperium by 50%

Child Targets

- Reduce under-five deaths due to HIV by at least 50%
- Provide antiretroviral therapy for all children with HIV

An estimated 7,000 children were newly infected with HIV in 2009. Modeling data from UNAIDS indicates that if all interventions are scaled up and Global Plan targets are achieved (see Figure 1), there would be 900 new child infections in 2015—an 87% decline in the number of new child infections between 2009 and 2015^[12].

Figure 1: Number of new child HIV infections due to mother to child transmission, by scenario, Côte d'Ivoire



Source:^[12] Joint United Nations Programme on HIV/AIDS, Unpublished estimates on PMTCT country targets, 2010

Global Plan Targets, Baseline and Current Status

	Indicators	2009 Baseline [or last available data]	2010	2011	2012	2013	2014	2015
Overall Targets	Number new paediatric HIV infections ^[3]	5,400	—	4,400	—	—	—	—
	Number HIV-associated maternal deaths ^[4]	1,400 (2005)	940	—	—	—	—	—
Child Targets	Percentage of under-5 deaths due to HIV	4% ^[13] (2008)	3% ^[7]	—	—	—	—	—
	ART coverage among children (ages 0-14) ^[3]	15%	—	15%	—	—	—	—
Prong One	HIV incidence in women (ages 15-49) ^[3]	0.19%	—	0.17%	—	—	—	—
Prong Two	Percentage of married women with unmet need for family planning (ages 15-49) ^[14]	29% (1999)	—	—	—	—	—	—
Prong Three	Mother-to-child transmission rate ^[3]	29%	—	27%	—	—	—	—
	Maternal ARV coverage (prophylaxis and ART coverage, excluding single-dose nevirapine) ^[3]	50% ^a	—	68%	—	—	—	—
	Breastfeeding ARV Coverage ^[3]	0%	—	0%	—	—	—	—
Prong Four	ART coverage among HIV+ pregnant women in need of treatment ^[3]	0%	—	0%	—	—	—	—

PRONG 1 & 2 Primary prevention of HIV among women of childbearing age Preventing unintended pregnancies among women living with HIV

Global 2015 Targets

- Reduce HIV incidence in women (ages 15-49) by 50%
- Reduce unmet need for family planning among women to zero

Background Statistics

Young people (ages 15-24) HIV prevalence _[1] *	Female: 1.5% [1.1-2.3%]	2009
	Male: 0.7% [0.5-1.1%]	2009
Condom use at last sex among young people (ages 15-24) with 2+ sexual partners in the last 12 months _[15]	Female: 45%	2005
	Male: 62%	2005
Male partners of pregnant women attending ANC tested in last 12 months _[9] *	Male: 3%	2010
Unintended pregnancies (ages 15-49)	Female: –	–

Key Points

Among young people (ages 15-24), HIV prevalence is two times higher among young women (1.5%) than young men (0.7%). Condom utilization at last sex among young people reporting multiple partners in the last year was higher among young men (62%) than young women (45%) in 2005, and coverage of HIV testing among male partners of pregnant women attending ANC is very low (3% in 2010).

PRONG 3 Preventing HIV transmission from a woman living with HIV to her infant


Global 2015 Targets

- Reduce overall mother-to-child transmission of HIV to <5%
- 90% of mothers receive perinatal ART or ARVs
- 90% of breastfeeding infant-mother pairs receive ART or ARVs

Background Statistics

Timing of 1 st ANC visit (months) _[15]	No ANC: 7%	2005
	<4 months: 31%	
	4-5 months: 33%	
	6-7 months: 22%	
	8+ months: 4%	
	DK: 4%	
Percentage of women attending at least 4 ANC visits during pregnancy _[15]	Total: 45%	2005
	Urban: 59%	
	Rural: 36%	
Percentage of pregnant women tested for HIV & received results in ANC, L&D, & post-partum (<72hrs) _[10] *	59%	2010
Estimated % of infants born to HIV+ women receiving ARVs for PMTCT _[10] *	44% [36-52%]	2010
Skilled attendant at delivery (%) _[16]	57%	2006
Exclusive breastfeeding for infants <6 months _[16]	4%	2006

Figure 2: Coverage of selected PMTCT interventions (2010)



† Indicator calculated from ANC health facility data for numerators and estimates of need for denominators. Not comparable to survey data presented under Background Statistics or elsewhere in this fact sheet.

Key Points

According to facility data, approximately 60% of pregnant women in Cote d'Ivoire attended at least one ANC visit in 2010, and 59% of pregnant women were tested for HIV. Some 68% of pregnant women living with HIV received more efficacious ARVs for PMTCT in 2011 and only 44% of HIV-exposed children received ARVs for PMTCT in 2010. According to 2006 data, only 57% of pregnant women access skilled attendance at delivery.

PRONG 4 Providing appropriate treatment, care and support to women living with HIV and their children and families

Global 2015 Target

• 90% of HIV-positive pregnant women in need of ART for their own health are started on lifelong ART

Background Statistics

Percentage of HIV-infected pregnant women assessed with CD4 testing ^[9] *	39%	2010
Percentage of infants born to HIV-infected women started on CTX prophylaxis within 2 months of birth ^[16] *	22% [18-26%]	2010
Percentage of infants born to HIV-infected women tested for HIV within 2 months of birth ^[10] *	36% [30-44%]	2010

Key Points

In 2010, only 39% of HIV-positive pregnant women were assessed with CD4 testing to determine if they were in need of ART for their own health. Coverage of CTX prophylaxis (22%) and early infant diagnosis (36%) within two months of birth was also low. Paediatric ART coverage has not changed since 2009 and remains very low—15% in 2011. Among pregnant women with HIV who received ARVs in 2010, 83% received more effective ARV regimens and 17% received ART for their own health.

Figure 3: Percentage distribution of various regimens provided to HIV+ pregnant women to prevent mother-to-child transmission of HIV, 2010^[9]*





Figure 4: Percentage of infants born to HIV+ women tested for HIV at 2 months of birth (EID) & Percentage of children living with HIV receiving ART, 2009-2011



Key Challenges & The Bottom Line

Key Challenges

Limited access to ANC and PMTCT services

Many women do not access skilled attendance at delivery

Many HIV+ pregnant women and HIV-exposed infants do not access ARVs for PMTCT

Few HIV+ pregnant women with HIV receive CD4 testing to determine if they are in need of treatment for their own health

Only a small proportion of infants born to HIV+ mothers receive CTX prophylaxis and are tested for HIV within two months of birth. Very few HIV+ children receive ART.

Data on some key PMTCT indicators are not available at national level, hindering tracking of progress towards the EMTCT targets

The Bottom Line

If national EMTCT targets for Côte d'Ivoire are to be met by 2015, the following actions should be considered:

Improve access to ANC and expand availability of PMTCT services within ANC to reach all women in need

Improve equitable access to skilled birth attendance

Improve access to more efficacious ARV regimens for HIV+ pregnant women and their children

Expand CD4 testing and enroll those eligible for treatment on ART

Expand the provision of CTX prophylaxis, EID, and ART for children with HIV

Strengthen routine monitoring system to ensure timely collection and reporting of quality data on Global Plan indicators.

References:

- 1 Joint United Nations Programme on HIV/AIDS, *Report on the Global AIDS Epidemic*, 2010
- 2 Joint United Nations Programme on HIV/AIDS, unpublished estimates, 2010
- 3 Joint United Nations Programme on HIV/AIDS, *Together We Will End AIDS*, 2012
- 4 World Health Organization, United Nations Children's Fund, United Nations Population Fund and the World Bank, *Trends in Maternal Mortality: 1990-2010*, Estimates developed by WHO, UNICEF, UNFPA and the World Bank, 2012
- 5 United Nations Children's Fund, *State of the World's Children 2012: Children in an Urban World, 2012*
- 6 United Nations Children's Fund, World Health Organization, the World Bank, United Nations DESA/Population Division, *Levels & Trends in Child Mortality, Report 2011*, Estimates Developed by the UN Inter-agency Group for Child Mortality Estimation, 2011
- 7 Liu L, Johnson HL, Cousens S, et al, for the Child Health Epidemiology Reference Group of WHO and UNICEF. *Global, regional, and national causes of child mortality: an updated systematic analysis for 2010 with time trends since 2000*. Lancet 2012
- 8 WHO/UNICEF/UNAIDS, calculated from Universal Access country reported unpublished data, 2011
- 9 World Health Organization, Joint United Nations Programme on HIV/AIDS, United Nations Children's Fund, *Towards Universal Access: Scaling up Priority HIV/AIDS Interventions in the Health Sector*. unpublished estimates, 2011
- 10 World Health Organization, Joint United Nations Programme on HIV/AIDS, United Nations Children's Fund, *Towards Universal Access: Scaling up Priority HIV/AIDS Interventions in the Health Sector. Progress report, 2011*
- 11 Republic of Côte d'Ivoire, Ministry of Health. *Plan national d'élimination de la transmission mère enfant du VIH à l'horizon 2015 au Côte d'Ivoire*
- 12 Joint United Nations Programme on HIV/AIDS, Unpublished estimates on PMTCT country targets, 2010
- 13 UN Inter-agency Group for Child Mortality Estimation, United Nations Children's Fund, World Health Organization, The World Bank, United Nations DESA/Population Division, *Levels & Trends in Child Mortality, Report 2010*, Estimates Developed by the UN Inter-agency Group for Child Mortality Estimation, UNICEF, WHO, The World Bank, United Nations DESA/Population Division, 2010
- 14 Côte d'Ivoire Demographic and Health Survey 1998-1999, Final Report

Notes:

- * Please note that the corresponding country data for this indicator have not been revised and, therefore, refer to what was published in the ***Global HIV/AIDS Response – Epidemic Update and Health Sector Progress Towards Universal Access, Progress Report 2011***. Revised country data for this indicator will be published towards the end of 2012.
- Data not available.
- † Indicator calculated from ANC health facility data for numerators and estimates of need for denominators. Not comparable to survey data presented under *Background Statistics* or elsewhere in this fact sheet.
- ^a 2009 data are not directly comparable to later years. Definition changed in 2010 to exclude single-dose nevirapine.

Acronyms:

- ANC: Antenatal care
- ART: Antiretroviral therapy
- ARVs: Antiretroviral prophylaxis
- CTX: Co-trimoxazole prophylaxis
- EID: Early infant diagnosis
- EMTCT: Elimination of mother-to-child transmission of HIV
- FP: Family planning
- L&D: Labour and delivery
- MMR: Maternal mortality ratio
- MNCH: Maternal, newborn and child health
- PMTCT: Prevention of mother-to-child transmission of HIV
- PWLHIV: Pregnant women living with HIV
- SRH: Sexual and reproductive health
- WHO: World Health Organization